

## Mouse Monoclonal Antibody to

# Amyloid $\beta$ A4 (40/42), C-Terminus

## clone 9F1

## FITC-labelled

**Order No.:** 0062-100FITC/bA4(40/42)-9F1

**Size ( $\mu$ g)** 100

**Lot No.:** 0062S



[www.nanotools.de](http://www.nanotools.de)

**orders & support:**

email: [info@nanotools.de](mailto:info@nanotools.de)

phone: +49-7641-455 670

fax: +49-7641-455 671

03/020307F

Isotype	Species	Reactivity	Applications	Mol. Weight	Ref.Cell Line	Epitope	Immunogen
IgG1	human		ICC		none	C-Terminus of Amyloid $\beta$ A4 (1-40) <u>and</u> $\beta$ A4 (1-42) crossreacts with APP	C-terminal peptide conjugated to KLH

### Background and Specificity:

The beta-amyloid peptide (beta A4), proteolytically released from the amyloid precursor protein (APP), is the principal component of senile plaques in Alzheimer's disease. Cleavage of APP by alpha-secretase or alternatively by beta-secretase leads to generation and extracellular release of soluble APP peptides, S-APP-alpha and S-APP-beta, respectively, and the retention of corresponding membrane-anchored C-terminal fragments, C83 and C99. Subsequent processing of C83 by gamma-secretase yields P3 peptides. This is the major secretory pathway and is nonamyloidogenic. Alternatively, presenilin/nicastrin-mediated gamma-secretase processing of C99 releases the amyloid beta proteins, amyloid-beta 40 (Abeta40) and amyloid-beta 42 (Abeta42), major components of amyloid plaques, and the cytotoxic C-terminal fragments, gamma-CTF(50), gamma-CTF(57) and gamma-CTF(59).

**Mab  $\beta$ A4(40/42)-9F1** interacts with the C-termini of both  $\beta$ -Amyloid (1 - 40) and (1-42).

### Related Products

**mab to  $\beta$ A4, N-Terminus**  
#0064-100/bA4N-19H5

**mab to  $\beta$ A4, N-Terminus**  
#0084-100/bA4N-19H11

**mab to  $\beta$ A4, N-Terminus**  
#0195-100/bA4N-7F4

**mab to  $\beta$ A4, N-Terminus**  
#0196-100/bA4N-7F9

**mab to  $\beta$ A4, N-Terminus**  
#0197-100/bA4N-11H3

**mab to  $\beta$ A4 (1-40), C-Terminus**  
#0060-100/bA4(40)-5C3

**mab to  $\beta$ A4 (1-40/42), C-Terminus**  
#0062-100/bA4(40/42)-9F1

**mab to  $\beta$ A4 (1-42), C-Terminus**  
#0061-100/bA4(42)-8G7

**mab to  $\beta$ A4 (1-43), C-Terminus**  
#0095-100/bA4(43)-6G12

<b>Purification:</b>	The antibody was purified from serum-free cell culture supernatant by subsequent thiophilic adsorption and size exclusion chromatography.
<b>Formulation:</b>	liquid; 0.5 mg/ml in PBS/0.09% Na-Azide/PEG and Sucrose
<b>Reconstitution:</b>	
<b>Stability:</b>	Aliquote and freeze in liquid nitrogen; store aliquots frozen at -80°C up to 1 year. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months.
	<b>Avoid repeated freeze / thaw cycles.</b>
<b>Positive Control:</b>	none
<b>Immunoblotting:</b>	Please use unlabelled or biotinylated antibody for this application. See related products.
<b>Immunoprecipitation:</b>	ND
<b>Immunocytochemistry:</b>	use at 1 - 10 $\mu$ g/ml
<b>ELISA:</b>	ND

All products are supplied for research and investigational use only. Not for use in humans or laboratory animals.